

# **Adeno-1 Expression System**

Enables efficient construction of recombinant adenoviruses.

Store kit at **-20°C**

Store Ad-competent cell at **-70°C**

## ***User Manual***

Version 2, January 2007

## 1. Introduction



**Application:** The Adeno-1 Expression System provides an efficient method for constructing recombinant adenoviruses. Our procedure uses *in vitro* ligation to subclone your gene of interest into a replication-incompetent (-E1/-E3) human adenoviral type 5 (Ad5) genome. This approach enables you to produce recombinant adenoviruses quickly (less than 3 weeks) and reliably. Our pShuttle vectors (+/-) have very flexible cloning sites, which allow easy subcloning manipulation.

## 2. Kit Contents

Table 1. Kit Contents.

Component	Quantity	Kit Verizon					
		v1.1	v1.2	v1.3	v1.4	v1.5	v1.6
Adeno DNA	5ug	✓	✓	✓			
pShuttle vector (+)	5ug		✓				
pShuttle vector (-)	5ug	✓		✓			
pShuttle-Laz control vector	5ug	✓	✓	✓			
Enzyme pack: PI-Sce I and I-CeuI; including buffer	2x50ul	✓			✓	✓	
Recombinant adenovirus PCR screening set	2x100ul	✓			✓	✓	
293 cells	1x10e6					✓	
Ad-competent cells	10x50ul						✓
Glycogen 20mg/ml	100ul						✓

- Store Ad-competent cells at -70°C and all other components at -20°C.
- Spin briefly to recover contents
- Avoid repeated freeze-thaw cycles



### 3. Experimental Procedure

#### 3.1 Cloning your gene of interest into the pShuttle (+/-):

3.1.1 Subclone your gene of interest into the pShuttle(+/-) using any of the unique restriction sites located in the MCS region.



## 3.2 Produce recombinant adenoviral DNA:

### 3.2.1 Subclone your expression cassette into the Adeno genome:

- a. Combine the reagents shown in Table 3 in a sterile 1.5 ml microcentrifuge tube in the order as shown.
- b. Gently mix, then spin briefly in the microcentrifuge.
- c. Incubate at room temperature for 1-2 hours.
- d. To the ligation product, add the following:
  - 1.5  $\mu$ l 10X Swa I digestion Buffer
  - 1.5  $\mu$ l 10X BSA
  - 1.0  $\mu$ l Swa I enzyme
  - 1.0  $\mu$ l ddH<sub>2</sub>O
- e. Incubate at room temperature for 1-2 hours.
- f. Perform phenol:chloroform:isoamyl alcohol (25:24:1) extraction by using the protocol in Appendix A.

**Table 3. Ligation of pShuttle to Adenovector**

Reagent	Volume
PI-Sce I/I-Ceu I digested vector	4ul
PI-Sce I/I-Ceu I digested pShuttle	3ul
5X DNA Ligation Buffer	2ul
DNA Ligase	1ul
<b>Total Volume</b>	<b>10ul</b>

### 3.2.2 Transform *E.coli* with ligation products

- a. Transform electro- or chemically competent DH<sub>5</sub> $\alpha$  or our **Ad-competent (Cat . VSA-0002)** cells with the SwaI digestion product.
- b. Select for ampicillin-resistant (Amp<sup>r</sup>) transformants by plating the transformation mixture on a LB agar/Amp plate (100  $\mu$ g/ml ampicillin). Incubate overnight at 37°C.
- c. Pick up 10 colonies and inoculate into 100  $\mu$ l LB/Amp medium in a sterile Eppendorf tube in the early morning.
- d. After 5-6 hours of incubation, screen for positive clones using PCR with Adeno screening primer mix (using the protocol in Appendix B).
- e. When you have identified a bacterial clone carrying the desired recombinant, inoculate 100-500 ml of LB/Amp

medium with all the bacteria (100 µl) from the 1.5 ml Eppendorf tube.

- f. Incubate the culture at 37°C overnight.
- g. Purify the plasmid using our *Column Pure Plasmid Maxi-Prep Kit (Cat No D205)* or the Qiagen kits. Expected yield: 30-50 µg plasmid DNA/100 ml culture.

**Note:** pAdeno is a large plasmid (>30 kb) that is susceptible to damage and rearrangement in E.coli. For best results, always use fresh, log-phase cultures for purification of recombinant pAdeno DNA. Do not store your culture at room temperature, 4°C, or on ice for long periods ( i.e. >24 hours) before starting purification.

### 3.2.3 Analyze putative recombinant Adenoviral DNA by PCR or restriction digestion

#### PI-Sce I and I-Ceu I Restriction Analysis:

- a. Set up 30µl PI-Sce I/I-Ceu I double digests by combining the reagents in Table 4 in a sterile 1.5 ml microcentrifuge tube.
- b. Mix well and spin briefly to collect contents.
- c. Incubate at 37°C for exactly 3 hours.
- d. Verify the insert by electrophoresis on a 0.8-1% agarose/*SafeView (Cat.No DG108)* gel.

**Important:** Since I-Ceu I and PI-Sce I tend to remain bound to DNA, use a gel loading buffer that contains SDS (final concentration after combining with sample: 0.1%), heat to 65°C for 10 minutes before loading.

**Table 4. Restriction Analysis of Recombinant Adenoviral DNA**

<b>Reagent</b>	<b>Volume</b>
Adenoviral DNA	3µg
10X Digestion Buffer	3ul
10X BSA	3ul
PI-Sce I (1 unit/ul)	2ul
I-Ceu I (1 unit/ul)	2ul
ddH2O	x ul
<b>Total Volume</b>	<b>30ul</b>

### PCR Analysis:

You can also screen pAdeno DNA for the presence of pShuttle derived expression cassettes by using PCR with Adeno forward and reverse PCR primers (using the protocol in Appendix B). These primers specifically amplify a 350-bp sequence that spans the cloning site in pAdeno. Only recombinant pAdenoviral templates are amplified since non-recombinants lack the pShuttle sequence needed for annealing with the reverse primer.

#### *3.2.4 Preparing Recombinant Adenoviral DNA for Transfection*

- a. In a sterile 1.5 ml microcentrifuge tube, combine the reagents in Table 5.
- b. Mix contents and spin the tube briefly.
- c. Incubate at 37°C for 2 hours
- d. Perform phenol extractions by using the protocol in Appendix A.

**Table 5. Pac I Digestion of Recombinant Adenoviral DNA**

<b>Reagent</b>	<b>Volume</b>
Recombinant Adenoviral DNA	3ug
10X Pac I Digestion Buffer	4ul
10X BSA	4ul
Pac I (10 units/ul)	2ul
ddH <sub>2</sub> O	x ul
<b>Total Volume</b>	<b>40ul</b>

### **3.3 Generate the recombinant adenovirus in packaging cells:**

1. Plate 293 cells at 70% confluency in a 6-well plate 12-24 hrs before transfection.
2. Incubate the plate(s) at 37°C in a humidified atmosphere maintained at 5% CO<sub>2</sub>.
3. Transfect each 6 well culture plate with 10 µl of Pac I-digested recombinant Adenoviral DNA. Use *Celfectin*<sup>TM</sup> (Cat.No G061) to transfer DNA into 293 cells.
4. Subculture cells into a T75 flask using trypsin 3-5 days after transfection.

5. Check periodically for cytopathic effect (CPE). **Note:** It normally takes 7-12 days to see CPE.
6. When >90% of the cells have detached from the plate, prepare viral stock by following steps 7-10. Name this stock “Primary Amplification” and store at –20°C.
  - Primary Amplification Stock is suitable for infecting target cells. We suggest you evaluate the function of this viral stock before preparing High-Titer Stock.
  - The presence of your recombinant construct can be verified by PCR or Western blotting.
7. Centrifuge the suspension at 1,500 x g for 5 min at room temperature.
8. Re-suspend the pellet in 500 µl sterile PBS
9. Lyse cells with three consecutive freeze-thaw-cycles. Freeze cells in dry ice/ethanol bath, thaw cells by placing the tube in a 37°C water bath. **Do not allow suspension to reach 37°C.**
10. After the third cycle, briefly centrifuge to pellet debris. Transfer the lysate to a clean, sterile centrifuge tube and store the lysate at –20°C.
11. Determine the adenoviral titer. The *Adenovector™ Rapid Titer Kit* (Cat.VSA-0024) enables you to determine the adenoviral titer by using an anti-Hexon antibody cell staining assay.

### **3.4 Infecting target cells:**

If the viruses are to be used in *in vitro* cell cultures, Double CsCl purification is not required as the viral supernatant will provide 100% gene transduction efficiency in most human cell lines. For *in vivo* studies (i.e. animal studies), purification is essential to remove defective particles, cell debris, and small amounts of media components, since these contaminants can induce significant immune responses. In addition, CsCl purification will concentrate the virus to a level suitable for *in vivo* injections.

## **Appendix A -Phenol:Chloroform:Isoamyl Alcohol Extraction**

1. Top up the sample to 100ul with TE buffer (pH 8.0), then add 100ul phenol:chloroform:isoamyl alcohol (25:24:1).
2. Vortex thoroughly.
3. Spin the tube in a microcentrifuge at 14,000 rpm for 2 min at room temperature to separate phases.
4. Carefully transfer the top aqueous layer to a clean 1.5-ml tube.
5. Add 400  $\mu$ l 95% ethanol, 25  $\mu$ l 10 M  $\text{NH}_4\text{OAc}$  (or 1/10 volume of 3 M  $\text{NaOAc}$ ), and 1  $\mu$ l glycogen (20 mg/ml).
6. Vortex thoroughly.
7. Spin the tube at 14,000 rpm for 5 min at room temperature.
8. Remove and discard the supernatant.
9. Carefully overlay the pellet with 300  $\mu$ l 70% ethanol.
10. Spin in a microcentrifuge at 14,000 rpm for 2 min at room temperature.
11. Carefully aspirate off the supernatant.
12. Air-dry the pellet for approximately 3-5 min at room temperature to evaporate residual ethanol.
13. Dissolve the DNA in 10  $\mu$ l sterile 1X TE Buffer (pH 8.0) for down-stream applications or store at  $-20^\circ\text{C}$  until use.

## **Appendix B -Identification of Recombinant Adenoviral DNA by PCR**

1. Set up the reaction as following using *2x PCR Taq MasterMix (Cat.No G013)* or your own Taq DNA polymerase:

Template (bacterial culture or purified DNA)	1 $\mu$ l
Primers (mix of both forward and reverse)	2 $\mu$ l
ddH <sub>2</sub> O	22 $\mu$ l
2x PCR Taq MasterMix	25 $\mu$ l
<b>Total</b>	<b>50 <math>\mu</math>l</b>

2. Perform PCR as following:

Denaturation	95°C	4 min
PCR Cycles (30-35)	94°C	30 sec
	55°C	30 sec
	72°C	1 min
	72°C	5 min
Final elongation	72°C	5 min

3. After PCR, perform DNA agarose gel electrophoresis. Positive band is at 350bp. Make sure to include a negative control for each assay.